Deterministic and Highly Efficient Quantum Cryptography with Entangled Photon Pairs

Zhi Zhao, Tao Yang, Zeng-Bing Chen, Jiangfeng Du, and Jian-Wei Pan
Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230027, People's Republic of China

We present a protocol for deterministic and highly efficient quantum cryptography with entangled photon pairs in a 4×4-dimentional Hilbert space. Two communicating parties, Alice and Bob first share a both polarization- and path-entangled photon pair, and then each performs a complete Bell-state measurement on their own photon in one of two complementary Bell-state bases. It is demonstrated that each measurement in which both Alice and Bob register a photon can build certain perfect correlation and generate 1.5 key bits on average. The security of our protocol is guaranteed by the non-cloning theorem.

PACS numbers: 03.67.Lx, 42.50.Ar

arXiv:quant-ph/0211098v2 11 Sep 2003

The central task for quantum cryptography is to establish secure keys before the transmission of message between two parties [1]. In recent years, both BB84 [2] and Ekert91 [3] cryptographic protocols have been successfully demonstrated in numerous experiments [4, 5, 6, 7, 8, 9]. However, one drawback of the above schemes is the non-deterministic feature, that is, only less than 50% of all detected qubits can be further used as key bits. This is a serious practical problem, because in the one-time-pad secret-key cryptosystem, the length of secret key must be the same as the ciphertext [10]. Although some deterministic cryptographic schemes based on orthogonal states have been proposed recently [11, 12, 13], the long storage rings required, which is an essential ingredient of those schemes, lead to a low efficiency for the transmission of polarized photons. More recently, deterministic quantum cryptography and secret direct communication have also been proposed with single-photon two-qubit states [14]. By exploiting the states in four- or more dimensional Hilbert space, each transmitted photon can be used as a key bit with the help of classical communication; eavesdroppers can be detected within the scheme. However, every photon detected can establish only one key bit.

In this work, we propose a deterministic and highly efficient quantum cryptography protocol using doubly entangled photon pairs [15, 16]. In the scheme, two communicating parties, Alice and Bob, first share a polarization- and path-entangled photon pair, then each performs a complete Bell measurement on Alice's (Bob's) photon in one of two complementary Bell bases. In this way, they can establish 1.5 key bits on average for each measurement where both Alice and Bob register a single photon. We also show that the security of our protocol is guaranteed by the non-cloning theorem.

In order to establish a key bit at each run - that is - in a determinstic way, it would require that, on the one hand, each assigned joint measurement must generate certain perfect correlations between two communicating parties; and on the other hand, some of these perfect correlations are sufficient to perform the validation of eavesdropper. Our quantum cryptography protocol is based on this observation. Although it is not possible to contruct such a protocol for two observers with only one Einstein-Poldolsky-Rosen (EPR) pair, one can indeed achieve this goal using two EPR pairs, provided that one can achieve the controlled-NOT operation between two indepedent photons [17].

In our scheme, instead of utilizing two EPR pairs, we make use of only one photon pair where the two photons are entangled both in polarization and in path degree of freedom [16]. In contrast to the two pair scheme, there are two obvious advantages in our single pair scheme. First, the required complete Bell-state measurement which projects Alice's (Bob's) photon onto a polarization-path entangled state in one of two complementary bases is feasible with only linear optics and, actually, has been experimentally implemented [18]. Second, using two usual polarization-entangled photon pairs will lead to very low efficiency due to the very low probability of simultaneously generating two entangled photon pairs. Our scheme is deterministic in the sense that all measurements in which both Alice and Bob register a photon can be used to establish perfect correlations, which can then be converted into key bits and a few of them are enough to perform the detection of any eavesdroppers. Entanglement exploited in the present scheme is used both to provide security against eavesdropping and to enhance the key bit generation.

We now explain how our scheme works. As shown in Fig. 1, two communicating parties, Alice and Bob, first share a both polarization- and path-entangled photon pair. It can be generated using parametric down conversion (PDC) by the method described in Ref. [19]. As explained in refs. [15, 16], after the UV laser passes through the BBO crystal twice, the state corresponding to the case where there is one and only one pair production is given by

$$|\Psi\rangle_{A1B2} = \frac{1}{2}\left(|H\rangle_1 |V\rangle_2 - |V\rangle_1 |H\rangle_2\right) \otimes \left(|a_1\rangle |b_2\rangle - |b_1\rangle |a_2\rangle\right). \quad (1)$$

$|H\rangle_1$ and $|V\rangle_1$ represent the polarization states of photon 1 in path a_1 or b_1, and $|H\rangle_2$ and $|V\rangle_2$ represent the states of photon 2 in path a_2 or b_2. Then let photon 1 in path a_1 and b_1 move towards Alice and photon 2 in path a_2 and b_2 towards Bob. With such state Alice and Bob can perform



FIG. 1: Schematic diagram of generating both polarization- and path-entangled photon pairs and performing the complete Bell measurement in one of the two complentary bases. R_{90} and R_{45} are half-wave plates that rotating the polarization of photons by 90^0 and 45^0 respectively. When inserting R_{90} into the mode a_1 (or b_2), Alice (or Bob) can perform the complete Bell-state measurement on the state of $\left\{\left|\Phi^{\pm}\right\rangle_{ij}, \left|\Psi^{\pm}\right\rangle_{ij}\right\}$. When inserting two 45^0s into the modes a_1 and b_1 (or a_2 and b_2) simultaneously, Alice (or Bob) can perform the complete Bell-state measurement on the state of $\left\{\left|\chi^{\pm}\right\rangle_{ij}, \left|\omega^{\pm}\right\rangle_{ij}\right\}$[18].

entanglement swapping [20, 21] or all-versus-nothing test of quantum mechanics against local realism [16, 17]. In the following, we are going to show how entanglement swapping of the quantum state $\left|\Psi\right\rangle_{A1B2}$ provides us a deterministic and efficient quantum cryptography.

In order to establish secure key bits between Alice and Bob, they can *randomly* choose one of two complementary Bell-state bases $\left\{\left|\Phi^{\pm}\right\rangle_{ij}, \left|\Psi^{\pm}\right\rangle_{ij}\right\}$ (type-I) or $\left\{\left|\chi^{\pm}\right\rangle_{ij}, \left|\omega^{\pm}\right\rangle_{ij}\right\}$ (type-II) $(ij = A1$ or $B2)$ and then perform Bell-state measurement. The two Bell-state bases are defined as following:

$$\begin{cases} \left|\Phi^{\pm}\right\rangle_{ij} = \frac{1}{\sqrt{2}}\left(\left|H\right\rangle_i \left|a_j\right\rangle \pm \left|V\right\rangle_i \left|b_j\right\rangle\right) \\ \left|\Psi^{\pm}\right\rangle_{ij} = \frac{1}{\sqrt{2}}\left(\left|H\right\rangle_i \left|b_j\right\rangle \pm \left|V\right\rangle_i \left|a_j\right\rangle\right) \end{cases}, \tag{2}$$

and

$$\begin{cases} \left|\chi^{\pm}\right\rangle_{ij} = \frac{1}{2}\left[\left|H\right\rangle_i \left(\left|a_j\right\rangle + \left|b_j\right\rangle\right) \pm \left|V\right\rangle_i \left(\left|a_j\right\rangle - \left|b_j\right\rangle\right)\right] \\ \left|\omega^{\pm}\right\rangle_{ij} = \frac{1}{2}\left[\left|V\right\rangle_i \left(\left|a_j\right\rangle + \left|b_j\right\rangle\right) \pm \left|H\right\rangle_i \left(\left|a_j\right\rangle - \left|b_j\right\rangle\right)\right] \end{cases}. \tag{3}$$

The Bell states in the two complementary bases satisfy the following relations

$$\begin{cases} \left|\Phi^{\pm}\right\rangle_{ij} = \frac{1}{\sqrt{2}}\left(\left|\chi^{\mp}\right\rangle_{ij} \pm \left|\omega^{\pm}\right\rangle_{ij}\right) \\ \left|\Psi^{\pm}\right\rangle_{ij} = \frac{1}{\sqrt{2}}\left(\left|\chi^{\pm}\right\rangle_{ij} \pm \left|\omega^{\mp}\right\rangle_{ij}\right) \end{cases}, \tag{4}$$

where $\left|\cdot\right\rangle_{A1}$ and $\left|\cdot\right\rangle_{B2}$ represent the quantum states to be measured by Alice and Bob, respectively. Because the Bell states are polarization and path entangled, the complete Bell-state measurement can be done with only linear optics [18]. This feature is essential to our deterministic quantum cryptography protocol. After the measurements have taken place, Alice and Bob can announce in public which type of Bell bases they have used, and then divide the measurements into two sub-groups: For the first group they use the same Bell bases, and for the second group they use the different Bell bases. Of course, owing to the limited detection efficiency they only consider those events in which both Alice and Bob successfully register a single photon.

Let us first consider what will happen when Alice and Bob use the same Bell bases, i.e., they use either type-I *or* type-II Bell base. In this case the state $\left|\Psi\right\rangle_{A1B2}$ of Alice and Bob, according to entanglement swapping, can be rewritten in the two Bell bases as

$$\begin{aligned} \left|\Psi\right\rangle_{A1B2} &= \frac{1}{2}\left(\left|\Phi^{+}\right\rangle_{A1}\left|\Phi^{+}\right\rangle_{B2} - \left|\Phi^{-}\right\rangle_{A1}\left|\Phi^{-}\right\rangle_{B2}\right. \\ &\quad \left. - \left|\Psi^{+}\right\rangle_{A1}\left|\Psi^{+}\right\rangle_{B2} + \left|\Psi^{-}\right\rangle_{A1}\left|\Psi^{-}\right\rangle_{B2}\right) \\ &= \frac{1}{2}\left(-\left|\chi^{+}\right\rangle_{A1}\left|\chi^{+}\right\rangle_{B2} + \left|\chi^{-}\right\rangle_{A1}\left|\chi^{-}\right\rangle_{B2}\right. \\ &\quad \left. + \left|\omega^{+}\right\rangle_{A1}\left|\omega^{+}\right\rangle_{B2} - \left|\omega^{-}\right\rangle_{A1}\left|\omega^{-}\right\rangle_{B2}\right). \end{aligned} \tag{5}$$

Eq. (5) implies that, whenever Alice obtain a measurement result, Bob will always obtain the same result as long as his measurement is performed in the same Bell bases. Therefore, perfect correlations can be built for those cases in which both of them perform a measurement on their own particle in the same Bell bases.

If Alice and Bob use different Bell bases (i.e., type-I *and* type-II Bell bases), the total state of Alice and Bob can be rewritten as either

$$\begin{aligned} \left|\Psi\right\rangle_{A1B2} &= \frac{1}{2\sqrt{2}}\left(\left|\Phi^{+}\right\rangle_{A1}\left|\omega^{+}\right\rangle_{B2} + \left|\Phi^{+}\right\rangle_{A1}\left|\chi^{-}\right\rangle_{B2}\right. \\ &\quad + \left|\Phi^{-}\right\rangle_{A1}\left|\omega^{-}\right\rangle_{B2} - \left|\Phi^{-}\right\rangle_{A1}\left|\chi^{+}\right\rangle_{B2} \\ &\quad - \left|\Psi^{+}\right\rangle_{A1}\left|\chi^{+}\right\rangle_{B2} - \left|\Psi^{+}\right\rangle_{A1}\left|\omega^{-}\right\rangle_{B2} \\ &\quad \left. + \left|\Psi^{-}\right\rangle_{A1}\left|\chi^{-}\right\rangle_{B2} - \left|\Psi^{-}\right\rangle_{A1}\left|\omega^{+}\right\rangle_{B2}\right), \end{aligned} \tag{6}$$

or

$$\begin{aligned} \left|\Psi\right\rangle_{A1B2} &= \frac{1}{2\sqrt{2}}\left(\left|\Phi^{+}\right\rangle_{B2}\left|\omega^{+}\right\rangle_{A1} + \left|\Phi^{+}\right\rangle_{B2}\left|\chi^{-}\right\rangle_{A1}\right. \\ &\quad + \left|\Phi^{-}\right\rangle_{B2}\left|\omega^{-}\right\rangle_{A1} - \left|\Phi^{-}\right\rangle_{B2}\left|\chi^{+}\right\rangle_{A1} \\ &\quad - \left|\Psi^{+}\right\rangle_{B2}\left|\chi^{+}\right\rangle_{A1} - \left|\Psi^{+}\right\rangle_{B2}\left|\omega^{-}\right\rangle_{A1} \\ &\quad \left. + \left|\Psi^{-}\right\rangle_{B2}\left|\chi^{-}\right\rangle_{A1} - \left|\Psi^{-}\right\rangle_{B2}\left|\omega^{+}\right\rangle_{A1}\right). \end{aligned} \tag{7}$$

The above two equations imply the following: Any specific outcome obtained by Alice or Bob in one of the two bases is maximally random and as such, given two specific outcomes of measurements in one type of the Bell bases obtained by Alice, she can predict with certainty the two outcomes of the corresponding measurements performed by Bob in another Bell bases and *vice versa*. For example, whenever Alice obtains one of the two Bell states $\left|\Phi^{+}\right\rangle$ and $\left|\Psi^{-}\right\rangle$ (or $\left|\Phi^{-}\right\rangle$ and $\left|\Psi^{+}\right\rangle$), Bob's photon will be in one of the two states $\left|\omega^{+}\right\rangle$ and $\left|\chi^{-}\right\rangle$ (or $\left|\omega^{-}\right\rangle$ and $\left|\chi^{+}\right\rangle$). Thus, Alice and Bob can always build two kinds of correlations in different bases.

Since a deterministic correlation can be built for every joint measurement, all outcomes of the measurements in the same or different bases can be further used as key bits.

This is the essential property of our deterministic quantum cryptography protocol and is different from the existing schemes with two-dimensional entangled photon pairs.

Having established that deterministic correlations can be built between two communicating parties, let us show how to generate the key bits in the present protocol. First, if Alice and Bob have used the different bases, they can then simply define the two kinds of correlations as 0 and 1. For example, in the case of that Alice performs a measurement in the $\left\{|\Phi^{\pm}\rangle_{ij}, |\Psi^{\pm}\rangle_{ij}\right\}$ bases and Bob performs a measurement in the $\left\{|\chi^{\pm}\rangle_{ij}, |\omega^{\pm}\rangle_{ij}\right\}$ bases, Alice can define the two Bell states $|\Phi^{+}\rangle$ and $|\Psi^{-}\rangle$ as 0 and $|\Phi^{-}\rangle$ and $|\Psi^{+}\rangle$ as 1, while Bob can define the two states $|\omega^{+}\rangle$ and $|\chi^{-}\rangle$ as 0 and $|\omega^{-}\rangle$ and $|\chi^{+}\rangle$ as 1. Thus in this case, one key bit can be built. Second, if they use the same Bell bases, they actually receive two bits of deterministic information from the measurements since there are 4 possible outcomes of measurements in any one of the two Bell bases. They can agree with each other in advance that the Bell states ($|\Phi^{+}\rangle$, $|\Phi^{-}\rangle$, $|\Psi^{+}\rangle$, $|\Psi^{-}\rangle$) and ($|\chi^{+}\rangle$, $|\chi^{-}\rangle$, $|\omega^{+}\rangle$, $|\omega^{-}\rangle$) are encoded as two bits 00, 01, 10 and 11. Thus in the latter case, two key bits can be built. Since the probabilities that Alice and Bob perform the mesaurement in different Bell bases and in the same Bell bases are the same, i.e. 50%, every measurement in which both of Alice and Bob register a photon can establish 1.5 key bits on average. Therefore, by exploiting the high-capacity encoding method, the above scheme can provide a more efficient quantum cryptography than the existing deterministic ones.

Let us now discuss the security of the scheme. For simplicity, suppose that an eavesdropper, Eve, has full access to the quantum channel between Alice and Bob, Eve can thus make use of intercept-resend attack. He can first randomly perform a Bell-state measurement on the photon 2 in the two complementary bases and then send its replacement to Bob. However, when Alice performs a Bell-state measurement on her photon by randomly choosing one of the two complementary bases, she equivalently prepares a mixture of some nonorthogonal states with equal probability. Owing to the limit of the non-cloning theorem [22], it is not possible for Eve to know which basis Alice has chosen. Indeed, Eve can at most get 50% information, while Alice and Bob have about 25% error rate in the key bits obtained [1, 2]. By announcing and comparing in public some outcomes of the measurements in the same bases, Alice and Bob can thus easily detect the presence of Eve.

However, Eve can also perform another type of intercept-resend attack. Eve can first randomly perform a Bell-state measurement on both photons in one of the two complementary bases and then send its replacements to Alice and Bob, respectively. In this case, since any measurement performed by Eve will destroy the original entanglement between the two photons, the measurement results obtained by Alice and Bob during the verifictaion of entanglement swapping will be inconsistent with the predictions of quantum mechanics. Specifically, by comparing the outcome of the measurement in the same Bell-state bases, in a single run Alice and Bob will detect the eavesdropper with the probability of $1/4$ whenever Eve performs the Bell measurements in the same Bell bases and with the probability of $1/2$ whenever Eve performs the Bell measurements in the different bases. By announcing and comparing in public some outcomes of the measurements in the same bases, again Alice and Bob can easily detect the presence of Eve.

As pointed out in Ref. [16], the state $|\Psi\rangle_{A1B2}$ used in our protocol is effectively a maximally entangled state in a 4×4-dimensional Hilbert space. Essentially, the scheme presented here is thus a high-dimensional quantum cryptography protocol. In contrast to the former high-dimensional quantum cryptography protocols where the protocols provide better security against eavesdropping than the low-dimensional ones while lead to a low generation rate of key bits [23, 24, 25, 26, 27], our protocol not only provides better security against eavesdropping but also has a higher generation rate of key bits than the low-dimensional ones.

Compared with Ekert's quantum cryptography protocol using EPR-entangled qubit states [3], a striking feature of the present scheme is its higher efficiency. In Ekert's scheme, in order to be against eavesdropping, Alice and Bob randomly perform statistical measurements and register the outcomes in one of three nonorthogonal bases. When they have used the same bases incidentally, the results are correlated or anticorrelated and can be converted into key bits. For the others the results is not correlated and therefore used only for checking whether there is eavesdropper with help of Bell's inequalities. Thus, the overall probability of creating a key bit is $2/9$ per pair. However, in our scheme each measurement in which both of Alice and Bob register a photon simultaneously can generate 1.5 key bits on average. Thus, the key bit generation efficiency per pair is $27/4$ times higher in our scheme than in Akert's.

In summary, we have presented a protocol for deterministic and efficient quantum cryptography with two photons that are entangled in both polarization and path degrees of freedom. The protocol can establish 1.5 key bits per entangled pair, which is almost one order of magnitude (~ 7 times) higher than the Akert's scheme and is completely secure against eavesdropping. The polarization- and path-entangled photon pairs [19] and the complete Bell-state measurement [18] are two basic ingredients of the proposed protocol. Thus, our scheme is feasible and within the reach of current technology. The experimental realization of the protocol is under way in our lab.

This work was supported by the National Natural Science Foundation of China, Chinese Academy of Sciences and the National Fundamental Research Program (under Grant No. 2001CB309303).

[1] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Rev. Mod. Phys. **74**, 145 (2002).
[2] C. H. Bennett and G. Brassard, in *Proceedings of the IEEE International Conference on Computers, Systems and Singal Proceeding,Bangalore, India* (IEEE, New York), p.175 (1984).
[3] A. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
[4] W. T. Butter *et al*., Phys. Rev. Lett. **81**, 3283 (1998).
[5] T. Jennewein *et al*., Phys. Rev. Lett. **84**, 4729 (2000).
[6] N. S. Naik *et al*., Phys. Rev. Lett. **84**, 4733 (2000).
[7] W. Tittel *et al*., Phys. Rev. Lett. **84**, 4737 (2000).
[8] W. T. Butter *et al*., Phys. Rev. Lett. **84**, 5652 (2000).
[9] C. Kurtsiefer *et al.*, Nature (London) **419**, 450 (2002).
[10] B. Schneier, *Applied Cryptography* (Wiley, New York, 1996).
[11] L. Goldenberg and L. Vaidman, Phys. Rev. Lett. **75**, 1239 (1995).
[12] M. Koashi and N. Imoto, Phys. Rev. Lett. **79**, 2383 (1997).
[13] A. Cabello, Phys. Rev. Lett. **85**, 5635 (2000).
[14] A. Beige, B. G. Englert, C. Kurtsiefer, and H. Weinfurter, Acta. Phys. Pol. A **101**, 357 (2002).
[15] C. Simon and J.-W. Pan, Phys. Rev. Lett. **89**, 257901 (2002).
[16] Z.-B. Chen, J.-W. Pan, Y.-D. Zhang, Č. Brukner, and A. Zeilinger, Phys. Rev. Lett. **90**, 160408 (2003).
[17] A. Cabello, Phys. Rev. Lett. **86**, 1911 (2001).
[18] D. Boschi *et al*., Phys. Rev. Lett. **80**, 1121 (1998).
[19] J.-W. Pan, S. Gasparoni, R. Ursin, G. Weihs, and A. Zeilinger, Nature **423**, 417 (2003).
[20] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993).
[21] J. -W. Pan, D. Bouwmeester, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. **80**, 3891 (1998).
[22] W. K. Wootters and W. H. Zurek, Nature (London) **299**, 802 (1982).
[23] D. Bruβ, Phys. Rev. Lett. **81**, 3018 (1998).
[24] H. Bechmann-Pasquinucci and N. Ginsin, Phys. Rev. A **59**, 4238 (1999).
[25] H. Bechmann-Pasquinucci and A. Peres, Phys. Rev. Lett. **85**, 3313 (2000).
[26] H. Bechmann-Pasquinucci and W. Tittel, Phys. Rev. A **61**, 062308 (2000).
[27] M. Bourennane, A. Karlsson, and G. Björk, Phys. Rev. A **64** 012306 (2001).